

EXISTENCE OF GOOD SWEEPOUTS ON CLOSED MANIFOLDS

LONGZHI LIN AND LU WANG

ABSTRACT. In this note we establish estimates for the harmonic map heat flow from S^1 into a closed manifold, and use it to construct sweepouts with the following good property: each curve in the tightened sweepout, whose energy is close to the maximal energy of curves in the sweepout, is itself close to a closed geodesic.

1. INTRODUCTION

Given a minimizing sequence of sweepouts of the width (see (3.1)), we apply the harmonic map heat flow on each curve in the sweepout to pull it tight while preserving the sweepout. Moreover the tightened sweepout has the following good property (see Theorem 3.4): each curve in the tightened sweepout whose energy is close to the maximal energy of curves in the sweepout is itself close to a closed geodesic. In particular, the width is the energy of some closed geodesic. On closed non-simply-connected surfaces, Grayson showed that there exist simple closed geodesics in each nontrivial π_1 homotopy class by the curve shortening flow; see [8]. On the 2-sphere whose π_1 homotopy group is trivial, Birkhoff used sweepouts to find non-trivial closed geodesics; see [1], [2], [6], [4], [3], [12] and section 2 in [7] about Birkhoff's idea. The argument works equally well on other Riemannian manifolds. The Birkhoff's curve shortening process is a discrete gradient flow of the length functional and the key point is a convexity formula for the energy functional which controls the distance of curves from closed geodesics explicitly; see Lemma 4.2 in [6] and cf. Theorem 3.1 in [5]. However, it requires some work to show the discrete shortening process preserves the homotopy class of sweepouts. Instead, we use a continuous method, i.e. the harmonic map heat flow, to tighten sweepouts, which provides a natural homotopy of sweepouts. There are several applications of the existence of good sweepouts on closed manifolds. For instance, in [6], Colding and Minicozzi bounded from above, by a negative constant, the rate of change of the width for a one-parameter family of convex hypersurfaces that flows by mean curvature. The estimate is sharp and leads to a sharp estimate for the extinction time. And a similar bound for the rate of change for the two dimensional width is shown for homotopy 3-spheres evolving by the Ricci flow; see [5] and [14].

Acknowledgements. We would like to thank our advisors Prof. Tobias Colding and Prof. William Minicozzi for their continuous guidance.

2. HARMONIC MAP HEAT FLOW

Throughout we use the subscripts θ and t to denote taking partial derivatives of maps with respect to θ and t ; u satisfies the harmonic map heat flow equation, which is defined in (2.1). Let (M, g) be a closed Riemannian manifold. By the

Nash Embedding Theorem, M can be isometrically embedded into Euclidean space $(\mathbb{R}^N, \langle \cdot, \cdot \rangle)$. Given a closed curve $\gamma \in H^1(S^1, M)$, define the energy functional $E(\gamma) = \frac{1}{2} \int_{S^1} |\gamma_\theta|^2 d\theta$. The harmonic map heat flow is the negative L^2 gradient flow of the energy functional. Thus the equation of the harmonic map heat flow from S^1 into M is

$$(2.1) \quad u_t = u_{\theta\theta} - A_u(u_\theta, u_\theta) \quad \text{on } (0, \infty) \times S^1; \quad \lim_{t \rightarrow 0+} u(t, \cdot) = u_0 \quad \text{in } H^1(S^1, M),$$

where A_u is the second fundamental form of M in $\mathbb{R}^N$ at point $u(\theta)$. We study the long time existence and uniqueness of the solution of (2.1). For relevant results of the two dimensional harmonic map heat flow, the reader could refer to [15]. Also we would like to thank Tobias Lamm for bringing to our attention the paper [13] of Ottarsson, which has some overlap with our paper and in which Theorem 2.1 was proved under the stronger assumption of C^1 initial data (and thus the C^1 continuity at $t = 0$)¹.

Theorem 2.1. *Given $u_0 \in H^1(S^1, M)$, there exists a unique solution $u(t, \theta) \in C^\infty((0, \infty) \times S^1, M)$ of (2.1).*

The following is devoted to the proof of Theorem 2.1. First, by the corollary on page 124 of [9], given any initial data $u_0 \in C^\infty(S^1, M)$, there exists $T_0 > 0$ and a unique solution $u \in C^\infty([0, T_0] \times S^1, M)$ of (2.1). We show that the solution u can be extended smoothly beyond T_0 . First, note that the energy is non-increasing under the harmonic map heat flow:

Lemma 2.2. *For $0 \leq t_1 \leq t_2 < T_0$,*

$$(2.2) \quad E(u(t_1, \cdot)) - E(u(t_2, \cdot)) = \int_{t_1}^{t_2} \int_{S^1} |u_t|^2 d\theta dt.$$

Proof. Multiply the harmonic map heat equation by u_t and integrate over $[t_1, t_2] \times S^1$,

$$\begin{aligned} \int_{t_1}^{t_2} \int_{S^1} |u_t|^2 d\theta dt &= \int_{t_1}^{t_2} \int_{S^1} \langle u_{\theta\theta}, u_t \rangle d\theta dt = - \int_{t_1}^{t_2} \int_{S^1} \langle u_\theta, u_{\theta t} \rangle d\theta dt \\ &= E(u(t_1, \cdot)) - E(u(t_2, \cdot)). \end{aligned}$$

□

Next we derive the gradient bound of u .

Lemma 2.3. $(\partial_t - \partial_\theta^2)|u_\theta|^2 \leq 0$.

Proof.

$$\begin{aligned} \partial_t |u_\theta|^2 &= 2\langle u_\theta, u_{\theta t} \rangle = 2\langle u_\theta, u_{\theta\theta\theta} \rangle - 2\langle u_\theta, (A_u(u_\theta, u_\theta))_\theta \rangle \\ &= 2\langle u_\theta, u_{\theta\theta\theta} \rangle + 2\langle u_{\theta\theta}, A_u(u_\theta, u_\theta) \rangle \\ &= \partial_\theta^2 |u_\theta|^2 - 2|u_{\theta\theta}|^2 + 2|A_u(u_\theta, u_\theta)|^2 \\ &\leq \partial_\theta^2 |u_\theta|^2. \end{aligned}$$

□

¹In our setting, the C^1 continuity at $t = 0$ may not be true. For our purpose that the harmonic map heat flow preserves the homotopy class of sweepouts, we use a different argument to show the H^1 continuity at $t = 0$.

Thus, by Theorem 7.36 in [11], for any $\tau > 0$ and $(t, \theta) \in [\tau, T_0] \times S^1$,

$$(2.3) \quad |u_\theta|^2(t, \theta) \leq \tau^{-1} E(u_0).$$

By Proposition 7.18 in [11], $|u_{\theta\theta}|$ and $|u_t|$ are bounded on $[2\tau, T_0] \times S^1$. And by induction, for any $\tau > 0$, the higher order derivatives of u on $[2\tau, T_0] \times S^1$ are bounded uniformly by constants depending only on M , $E(u_0)$, τ and T_0 . Hence u can be extended smoothly to a solution of (2.1) beyond T_0 . In other word, there exists a unique solution $u \in C^\infty([0, \infty) \times S^1, M)$ of (2.1), if $u_0 \in C^\infty(S^1, M)$.

Next, in general, given $u_0 \in H^1(S^1, M)$, we can find a sequence $u_0^m \in C^\infty(S^1, M)$ approaching u_0 in the H^1 topology. Let u^m be the solution of the harmonic map heat flow with initial data u_0^m . Thus by (2.3) and discussion above, for any $\tau > 0$ and $T_0 > \tau$, u^m and all their derivatives are bounded uniformly, independent of m . Hence by the Arzela-Ascoli Theorem and a diagonalization argument, there exists a map $u \in C^\infty((0, \infty) \times S^1, M)$ solving the harmonic map heat flow with $E(u(t, \cdot)) \leq E(u_0)$. And it follows from the lemma below that $t \rightarrow u(t, \cdot)$ is a continuous map from $[0, \infty) \rightarrow H^1(S^1, M)$.

Lemma 2.4. *Given $\epsilon > 0$, there exists $\delta > 0$, depending on M , u_0 and ϵ , so that if $0 \leq t_1 < t_2$ and $t_2 - t_1 < \delta$, then*

$$(2.4) \quad \|u(t_2, \cdot) - u(t_1, \cdot)\|_{H^1(S^1)} \leq \epsilon.$$

Proof. Note that by Lemma 2.2, $\lim_{t \rightarrow 0} u(t, \cdot) = u_0$ in the $L^2(S^1, M)$ topology. Moreover, we have

$$\int_{S^1} |u(t_2, \theta) - u(t_1, \theta)|^2 d\theta \leq \int_{S^1} \left| \int_{t_1}^{t_2} u_t dt \right|^2 d\theta \leq (t_2 - t_1) \int_{t_1}^{t_2} \int_{S^1} |u_t|^2 d\theta dt.$$

Next, by Lemma 2.2 and the Cauchy-Schwarz inequality,

$$\begin{aligned} & \int_{S^1} |u_\theta(t_2, \theta) - u_\theta(t_1, \theta)|^2 d\theta \\ &= \int_{S^1} |u_\theta(t_1, \theta)|^2 d\theta - \int_{S^1} |u_\theta(t_2, \theta)|^2 d\theta - 2 \int_{S^1} \langle u_\theta(t_2, \theta), u_\theta(t_1, \theta) - u_\theta(t_2, \theta) \rangle d\theta \\ &= 2 \int_{t_1}^{t_2} \int_{S^1} |u_t|^2 d\theta dt + 2 \int_{S^1} \langle u_{\theta\theta}(t_2, \theta), u(t_1, \theta) - u(t_2, \theta) \rangle d\theta \\ &\leq 2 \int_{t_1}^{t_2} \int_{S^1} |u_t|^2 d\theta dt + 2 \left(\int_{S^1} |u_{\theta\theta}(t_2, \theta)|^2 d\theta \right)^{\frac{1}{2}} \left(\int_{S^1} |u(t_2, \theta) - u(t_1, \theta)|^2 d\theta \right)^{\frac{1}{2}} \\ &\leq 2 \int_{t_1}^{t_2} \int_{S^1} |u_t|^2 d\theta dt + 2(t_2 - t_1)^{\frac{1}{2}} \left(\int_{S^1} |u_{\theta\theta}(t_2, \theta)|^2 d\theta \right)^{\frac{1}{2}} \left(\int_{t_1}^{t_2} \int_{S^1} |u_t|^2 d\theta dt \right)^{\frac{1}{2}}. \end{aligned}$$

From Lemma 2.2, (2.1) and (2.3),

$$(2.5) \quad \int_{S^1} |u_{\theta\theta}(t_2, \theta)|^2 d\theta \leq \int_{S^1} |u_t(t_2, \theta)|^2 d\theta + t_2^{-1} \sup_M |A|^2 \cdot E(u_0)^2.$$

We derive the evolution equation of $|u_t|^2$:

$$\begin{aligned} \partial_t |u_t|^2 &= 2 \langle u_t, u_{tt} \rangle = 2 \langle u_t, u_{\theta\theta t} \rangle - 2 \langle u_t, (A_u(u_\theta, u_\theta))_t \rangle \\ &= \partial_\theta^2 |u_t|^2 - 2 |u_{t\theta}|^2 + 2 \langle u_{tt}, A_u(u_\theta, u_\theta) \rangle \\ &= \partial_\theta^2 |u_t|^2 - 2 |u_{t\theta}|^2 + 2 \langle A_u(u_t, u_t), A_u(u_\theta, u_\theta) \rangle. \end{aligned}$$

Thus by (2.3), if $t > t_3 = t_1 + (t_2 - t_1)/2$,

$$(2.6) \quad (\partial_t - \partial_\theta^2)|u_t|^2 - 4(t_2 - t_1)^{-1} \sup_M |A|^2 \cdot E(u_0) \cdot |u_t|^2 \leq 0.$$

Hence

$$\begin{aligned} \int_{S^1} |u_t|^2(t_2, \theta) d\theta &\leq \inf_{t_3 \leq t \leq t_2} \int_{S^1} |u_t|^2(t, \theta) d\theta + C(t_2 - t_1)^{-1} \int_{t_1}^{t_2} \int_{S^1} |u_t|^2 d\theta dt \\ &\leq (C + 2)(t_2 - t_1)^{-1} \int_{t_1}^{t_2} \int_{S^1} |u_t|^2 d\theta dt, \end{aligned}$$

where C depends on M and $E(u_0)$. Combining the inequality above, (2.5) and Lemma 2.2, there exists $\delta > 0$ so that (2.4) holds. $\square$

It follows from Lemma 2.4 and (2.3) that there exists $R_0 > 0$, depending only on M and u_0 , so that for $t \geq 0$, $\sqrt{2\pi} \cdot \sup_M |A|^2 \cdot \int_{\{t\} \times I_{R_0}} |u_\theta|^2 d\theta < 1/16$, where I_{R_0} is any segment on unit circle of length R_0 . To prove the uniqueness of the solution of (2.1), we need the following lemma.

Lemma 2.5. *Suppose that u is a solution of (2.1) in $C^\infty((0, \infty) \times S^1, M)$. Then*

$$(2.7) \quad \int_0^T \int_{S^1} |u_{\theta\theta}|^2 d\theta dt \leq \frac{T}{4R_0^2} E(u_0) + 2[E(u_0) - E(u(T, \cdot))].$$

Proof. If $0 < t_0 \leq T$, then by (2.1) and an interpolation inequality (see Lemma 6.7 in [16]),

$$\begin{aligned} \int_{t_0}^T \int_{S^1} |u_{\theta\theta}|^2 d\theta dt &\leq \int_{t_0}^T \int_{S^1} |u_t|^2 d\theta dt + \sup_M |A|^2 \cdot \int_{t_0}^T \int_{S^1} |u_\theta|^4 d\theta dt \\ &\leq [E(u_0) - E(u(T, \cdot))] + \frac{1}{2} \int_{t_0}^T \int_{S^1} |u_{\theta\theta}|^2 d\theta dt + \frac{T}{8R_0^2} E(u_0). \end{aligned}$$

Absorbing the righthand side into the lefthand side and noting that the estimate is independent of t_0 , (2.7) follows immediately. $\square$

Now we are ready to show the uniqueness of the solution of the harmonic map heat flow.

Lemma 2.6. *Given $u_0 \in H^1(S^1, M)$, let u and $\tilde{u}$ be solutions of (2.1) in $C^\infty((0, \infty) \times S^1, M)$. Then $u = \tilde{u}$.*

Proof. Define $v = u - \tilde{u}$, thus

$$(2.8) \quad v_t = v_{\theta\theta} - A_u(u_\theta, u_\theta) + A_{\tilde{u}}(\tilde{u}_\theta, \tilde{u}_\theta).$$

Multiply both sides of (2.8) by v and integrate over $[0, t_0] \times S^1$,

$$\begin{aligned}
& \int_{\{t_0\} \times S^1} |v|^2 d\theta + 2 \int_0^{t_0} \int_{S^1} |v_\theta|^2 d\theta dt \\
&= 2 \int_0^{t_0} \int_{S^1} \langle A_{\tilde{u}}(\tilde{u}_\theta, \tilde{u}_\theta) - A_u(u_\theta, u_\theta), v \rangle d\theta dt \\
&\leq C(M) \int_0^{t_0} \int_{S^1} |v|^2 (|\tilde{u}_\theta|^2 + |u_\theta|^2) d\theta dt + C(M) \int_0^{t_0} \int_{S^1} |v| |v_\theta| (|\tilde{u}_\theta| + |u_\theta|) d\theta dt \\
&\leq C(M) \int_0^{t_0} \int_{S^1} |v|^2 (|\tilde{u}_\theta|^2 + |u_\theta|^2) d\theta dt + \int_0^{t_0} \int_{S^1} |v_\theta|^2 d\theta dt \\
&\leq C(M) \int_0^{t_0} \left(\|u_\theta\|_{C^0(\{t\} \times S^1)}^2 + \|\tilde{u}_\theta\|_{C^0(\{t\} \times S^1)}^2 \right) \int_{S^1} |v|^2 d\theta dt + \int_0^{t_0} \int_{S^1} |v_\theta|^2 d\theta dt.
\end{aligned}$$

By Lemma 2.2, 2.5 and Sobolev embedding theorem, there exists $\delta > 0$, depending on M and u_0 , so that if $t_0 \leq \delta$, then

$$\begin{aligned}
& C(M) \int_0^{t_0} \|\tilde{u}_\theta\|_{C^0(\{t\} \times S^1)}^2 + \|u_\theta\|_{C^0(\{t\} \times S^1)}^2 dt \\
&\leq C(M) \int_0^{t_0} \int_{S^1} |\tilde{u}_{\theta\theta}|^2 + |u_{\theta\theta}|^2 d\theta dt \\
&\leq C(M) \left[E(u_0) \frac{t_0}{2R_0^2} + 4E(u_0) - 2E(u(t_0, \cdot)) - 2E(\tilde{u}(t_0, \cdot)) \right] \leq \frac{1}{2}.
\end{aligned}$$

Thus, absorbing the righthand side into the lefthand side,

$$(2.9) \quad \sup_{0 \leq t \leq \delta} \int_{\{t\} \times S^1} |v|^2 d\theta + 2 \int_0^\delta \int_{S^1} |v_\theta|^2 d\theta dt \leq 0.$$

Since $[0, T]$ is compact, Lemma 2.6 follows by iteration. $\square$

3. WIDTH AND GOOD SWEEPOUTS

In [6], Colding and Minicozzi introduced the crucial geometric concepts: sweepout and width.

Definition 3.1. A continuous map $\sigma : [-1, 1] \times S^1 \longrightarrow M$ is called a sweepout on M , if $\sigma(s, \cdot) \in H^1(S^1, M)$ for each $s \in [-1, 1]$, the map $s \longrightarrow \sigma(s, \cdot)$ is continuous from $[-1, 1]$ to $H^1(S^1, M)$ and σ maps $\{-1\} \times S^1$ and $\{1\} \times S^1$ to points.

The sweepout σ induces a map $\tilde{\sigma}$ from the sphere S^2 to M and we will not distinguish σ with $\tilde{\sigma}$. Denote by Ω the set of sweepouts on M . The homotopy class Ω_σ of σ is the path connected component of σ in Ω , where the topology is induced from $C^0([-1, 1], H^1(S^1, M))$.

Definition 3.2. The width $W = W(\Omega_\sigma)$ of the homotopy class Ω_σ is defined by

$$(3.1) \quad W = \inf_{\tilde{\sigma} \in \Omega_\sigma} \max_{s \in [-1, 1]} E(\tilde{\sigma}(s, \cdot)).$$

For $\alpha \in (0, 1)$ fixed, let $\gamma : S^1 \longrightarrow M$ be a smooth closed curve and G be the set of closed geodesics in M . Define $\text{dist}_\alpha(\gamma, G) = \inf_{\tilde{\gamma} \in G} \|\gamma - \tilde{\gamma}\|_{C^{1, \alpha}(S^1)}$. We prove the following proposition for the solution of (2.1), which is the key to the proof of Theorem 3.4:

Proposition 3.3. *Given $0 < \alpha < 1$, $W_0 \geq 0$, $t_0 > 0$ and $\epsilon > 0$, there exists $\delta_0 > 0$ so that if $W_0 - \delta_0 \leq E(u(t_0, \cdot)) \leq E(u_0) \leq W_0 + \delta_0$, then $\text{dist}_\alpha(u(t_0, \cdot), G) < \epsilon$.*

Proof. If Proposition 3.3 fails, then there exist $0 < \alpha < 1$, $W_0 \geq 0$, $t_0 > 0$, $\epsilon > 0$, and a sequence of solutions u^j of the harmonic map heat flow satisfying that $W_0 - 1/j \leq E(u^j(t_0, \cdot)) \leq E(u_0^j) \leq W_0 + 1/j$ and $\text{dist}_\alpha(u^j(t_0, \cdot), G) \geq \epsilon$. It follows from the evolution equation of $|u_t^j|^2$ (see (2.6)), Lemma 2.2 and Theorem 7.36 in [11] that

$$(3.2) \quad \sup_{\theta \in S^1} |u_t^j|^2(t_0, \theta) \leq C [E(u^j(t_0/2, \cdot)) - E(u^j(t_0, \cdot))],$$

where C depends on M , t_0 and W_0 . Thus $\sup_{\theta \in S^1} |u_t^j|(t_0, \theta) \rightarrow 0$ and it follows from (2.3) that $\|u^j(t_0, \cdot)\|_{C^2(S^1)}$ is uniformly bounded by constants depending on M , t_0 and W_0 . Therefore by Arzela-Ascoli's Theorem and Theorem 1.5.1 in [10], there exists a subsequence (relabelled) $u^j(t_0, \cdot)$ converges to u^∞ in $C^{1,\alpha}(S^1, M)$ and u^∞ is a closed geodesic in M . This is a contradiction. $\square$

Let σ be a sweepout on closed manifold M and σ^j be a minimizing sequence of sweepouts in Ω_σ , that is

$$(3.3) \quad W \leq \max_{s \in [-1, 1]} E(\sigma^j(s, \cdot)) \leq W + 1/j.$$

Applying the harmonic map heat flow to each slice of σ^j , we get a map $\Phi^j : [-1, 1] \times [0, \infty) \times S^1 \rightarrow M$ and for each $s \in [-1, 1]$ fixed, $\Phi^j(s, t, \theta)$ solves (2.1) with $\Phi^j(s, 0, \theta) = \sigma^j(s, \theta)$. It follows from the proof of the long time existence and uniqueness of the solution of (2.1) that for any $t_0 \geq 0$, the map $s \rightarrow \Phi(s, t_0, \cdot)$ is continuous from $[-1, 1]$ to $H^1(S^1, M)$ and thus $\Phi^j(\cdot, t_0, \cdot)$ is still a sweepout on M . Since $[-1, 1]$ is compact, for any $\epsilon > 0$, there exists $\delta > 0$ so that: if $0 \leq t_1 < t_2 \leq t_0$ and $t_2 - t_1 < \delta$, then $\int_{t_1}^{t_2} \int_{S^1} |\Phi_t^j(s, t, \theta)|^2 d\theta dt < \epsilon$ for any $s \in [-1, 1]$. Hence by Lemma 2.4, for any $t_0 > 0$, $\Phi^j(\cdot, t_0, \cdot)$ is homotopic to σ^j . Therefore it follows from Proposition 3.3 that the $\Phi^j(\cdot, t_0, \cdot)$ are good sweepouts on M . That is,

Theorem 3.4. *Given $0 < \alpha < 1$, $t_0 > 0$ and $\epsilon > 0$, there exists $\delta > 0$ so that if $j > 1/\delta$ and $s \in [-1, 1]$ satisfies $E(\Phi^j(s, t_0, \cdot)) \geq W - \delta^2$, then $\text{dist}_\alpha(\Phi^j(s, t_0, \cdot), G) < \epsilon$.*

In [6], Colding and Minicozzi show that the width is non-negative and is positive if σ is not the zero element in $\pi_2(M)$. In fact, assume that $W(\Omega_\sigma) = 0$ and $\hat{\sigma} \in \Omega_\sigma$ is such that the energy of each slice of $\hat{\sigma}$ is sufficiently small. Then each slice, $\hat{\sigma}(s, \cdot)$, is contained in a strictly convex neighborhood of $\hat{\sigma}(s, \theta_0)$ and note that $s \rightarrow \hat{\sigma}(s, \theta_0)$ is a continuous curve on M . Hence a geodesic homotopy connects $\hat{\sigma}$ to a path of point curves and thus $\hat{\sigma}$ is homotopically trivial. Since G is closed in the $H^1(S^1, M)$ topology, we have

Corollary 3.5. *If M is a closed Riemannian manifold and $\pi_2(M) \neq \{0\}$, then there exists at least one non-trivial closed geodesic on M .*

Remark 3.6. Instead of using the unit interval $[-1, 1]$ as the parameter space for the curves in the sweepout and assuming that the curves start and end in point curves, we could have used any compact space $\mathcal{P}$ and required that the curves are constants on $\partial\mathcal{P}$. In this case, $\Omega^\mathcal{P}$ is the set of continuous maps $\sigma : \mathcal{P} \times S^1 \rightarrow M$ so that for each $s \in \mathcal{P}$ the curve $\sigma(s, \cdot)$ is in $H^1(S^1, M)$, the map $s \rightarrow \sigma(s, \cdot)$ is

²Such s exists, since $W \leq \max_{s \in [-1, 1]} E(\Phi^j(s, t_0, \cdot)) \leq W + 1/j$.

continuous from $\mathcal{P}$ to $H^1(S^1, M)$, and finally σ maps $\partial\mathcal{P}$ to point curves. Given $\sigma \in \Omega^{\mathcal{P}}$, the homotopy class $\Omega_{\sigma}^{\mathcal{P}}$ is the set of maps $\hat{\sigma} \in \Omega^{\mathcal{P}}$ that are homotopic to σ through maps in $\Omega^{\mathcal{P}}$. And the width $W = W(\Omega_{\sigma}^{\mathcal{P}})$ is defined by

$$(3.4) \quad W = \inf_{\hat{\sigma} \in \Omega_{\sigma}^{\mathcal{P}}} \max_{s \in \mathcal{P}} E(\hat{\sigma}(s, \cdot)).$$

Theorem 3.4 holds for general parameter space; the proof is virtually the same when $\mathcal{P} = [-1, 1]$.

REFERENCES

- [1] G.D. Birkhoff. Dynamical systems with two degrees of freedom. *TAMS*, 18(2):199–300, 1917.
- [2] G.D. Birkhoff. Dynamical systems, volume 9. AMS Colloq. Publ., Providence, RI, 1927.
- [3] T.H. Colding and C. De Lellis. The min-max construction of minimal surfaces. In *Surveys in differential geometry*, Vol.8 (Boston, MA, 2002), pages 75–107. Int. Press, Somerville, MA, 2003.
- [4] T.H. Colding and W.P. Minicozzi. Estimates for the extinction time for the Ricci flow on certain 3-manifolds and a question of Perelman. *J. Amer. Math. Soc.*, 18(3):561–569, 2005.
- [5] T.H. Colding and W.P. Minicozzi. Width and finite extinction time of Ricci flow. *Geom. Topol.*, 12(5):2537–2586, 2008.
- [6] T.H. Colding and W.P. Minicozzi. Width and mean curvature flow. *Geom. Topol.*, 12(5):2517–2535, 2008.
- [7] C.B. Croke. Area and the length of the shortest closed geodesic. *J. Diff. Geom.*, 27(1):1–21, 1988.
- [8] M.A. Grayson. Shortening embedded curves. *Ann. of Math. (2)*, 129(1):71–111, 1989.
- [9] R.S. Hamilton. Harmonic maps of manifolds with boundary, volume 471 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1975.
- [10] F. Hélein. Harmonic maps, conservation laws and moving frames, volume 150 of *Cambridge Tracts in Mathematics*. Cambridge University Press, second edition, 2002.
- [11] G.M. Lieberman. Second order parabolic differential equations. World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [12] L. Lin. Closed geodesics in Alexandrov spaces of curvature bounded from above. *preprint*.
- [13] S.K. Ottarsson. Closed geodesics on Riemannian manifolds via the heat flow. *J. Geom. Phys.*, 2(1):49–72, 1985.
- [14] G. Perelman. Finite extinction time for the solutions to the Ricci flow on certain three manifolds. *math.DG/0307245*.
- [15] M. Struwe. On the evolution of harmonic mappings of Riemannian surfaces. *Comment. Math. Helv.*, 60(4):558–581, 1985.
- [16] M. Struwe. Variational methods. Applications to nonlinear partial differential equations and Hamiltonian systems, volume 34 of *Ergebnisse der Mathematik und ihrer Grenzgebiete 3. Folge. Band 34. A Series of Modern Surveys in Mathematics*. Springer-Verlag, Berlin, second edition, 2002.

DEPARTMENT OF MATHEMATICS, JOHNS HOPKINS UNIVERSITY, 3400 N. CHARLES ST., BALTIMORE, MD 21218

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, 77 MASSACHUSETTS AVENUE, CAMBRIDGE, MA 02139

E-mail address: lzlin@math.jhu.edu and luwang@math.mit.edu